

## University of Richmond UR Scholarship Repository

---

Master's Theses

Student Research

---

1974

# In vitro effects of prolactin, cortisol, aldosterone, and cyclic amp on branchial Na<sup>+</sup>, K<sup>+</sup> - activated ATPase of the killifish, *Fundulus heteroclitus*

Stephen Burke Kerby

Follow this and additional works at: <http://scholarship.richmond.edu/masters-theses>

---

### Recommended Citation

Kerby, Stephen Burke, "In vitro effects of prolactin, cortisol, aldosterone, and cyclic amp on branchial Na<sup>+</sup>, K<sup>+</sup> - activated ATPase of the killifish, *Fundulus heteroclitus*" (1974). *Master's Theses*. Paper 373.

This Thesis is brought to you for free and open access by the Student Research at UR Scholarship Repository. It has been accepted for inclusion in Master's Theses by an authorized administrator of UR Scholarship Repository. For more information, please contact [scholarshiprepository@richmond.edu](mailto:scholarshiprepository@richmond.edu).

IN VITRO EFFECTS OF PROLACTIN, CORTISOL, ALDOSTERONE,  
AND CYCLIC AMP ON BRANCHIAL  $\text{Na}^+$ ,  $\text{K}^+$  - ACTIVATED ATPASE  
OF THE KILLIFISH, FUNDULUS HETEROCLITUS

by

STEPHEN BURKE KERBY

B.S., GEORGE MASON UNIVERSITY

Approved:

David W. Toule  
Committee Chairman

Austin D. Smith  
Dean of the Graduate School

Examining Committee:

F. B. Leftwich

W. R. Gennery

R. D. Decker

\_\_\_\_\_

\_\_\_\_\_

B. C. Schaefer

M. E. Gennery

H. E. Rice

\_\_\_\_\_

\_\_\_\_\_

## TABLE OF CONTENTS

Acknowledgements .....	11
Abstract .....	111
Introduction .....	1
Methods and Materials .....	10
Results and Discussion .....	13
Literature Cited .....	17
Tables .....	28
Figures .....	32
Vita .....	33

## ACKNOWLEDGEMENTS

I would like to express my sincere appreciation to Dr. David W. Towle, my thesis advisor, and to Drs. Francis B. Leftwich and Wilton R. Tenney, my reading committee members, for their aid in the preparation of this thesis.

I especially thank Ms. Barbara Phillips for her help in typing this thesis.

# ABSTRACT

In vitro effects of prolactin, cortisol, aldosterone, and cyclic adenosine monophosphate were studied on branchial  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase of freshwater adapted killifish, Fundulus heteroclitus.

Decreases in ATPase activity were found after treatment with prolactin and aldosterone. Cyclic AMP caused an increase in ATPase activity.

The action of cortisol on this enzyme was not clear. A dose - response relationship was found only for cyclic AMP. It is proposed that cyclic AMP may be involved in the adaptation of F. heteroclitus to seawater.

## INTRODUCTION

The comparative physiology of salt and water homeostasis in fishes has long fascinated biologists interested in the mechanism of sodium transport, because the demands of the external environment are so stringent, and the evolutionary adjustments arrived at are so varied (Parry, 1966; Smith, 1930). The problem facing animal cells is that sodium enters the cell by diffusion (a passive process) with a concentration gradient and then must be extruded against that gradient (an active process). Potassium leaks from the cell with a concentration gradient and then must be actively reabsorbed (Fraser, 1970). This active process, known as the sodium pump, requires energy which is considered to be derived from a labile high energy phosphate group present in adenosine triphosphate (ATP), and which becomes available when ATP is broken down to adenosine diphosphate (Bentley, 1971). The release of energy from ATP and its use to drive the sodium pump is thought to be catalyzed by a specific adenosine triphosphatase (ATPase), activated by sodium and potassium (Skou, 1965). Sodium - potassium - activated adenosine triphosphatase ( $\text{Na}^+$ ,  $\text{K}^+$  - ATPase), first isolated from invertebrate nervous tissue by Skou in 1957, plays a key role in the active reciprocal transfer of sodium and potassium

across the plasma membrane of individual cells. There is increasing evidence that this enzyme is intimately involved in the secretion of sodium across teleost epithelial membranes in the kidney (Katz and Epstein, 1967; Pickford et al., 1970), the urinary bladder (Hirano et al., 1971; Lam, 1972), the intestine (Utida et al., 1972; Lam, 1972), and the gill (Epstein et al., 1967; Kamiya and Utida, 1969; Milne et al., 1971). The specific activity of  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase in these tissues appears to be proportional to the level of sodium transport or pumping activity that the tissue would be expected to carry out in response to the osmotic stress placed on the animal by the environment (Jampol and Epstein, 1970).

Freshwater teleosts maintain a serum sodium level of about 140 - 160 mEq / liter by reducing the efflux of sodium to a minimum through the gill and kidney, by active inward transport of a small amount of sodium by the gills, and by excreting large quantities of water through the kidneys (Maetz, 1964; Parry, 1966; Potts, 1968; Jampol and Epstein, 1970). Consistent with these ion transport requirements,  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity is low in the gills and high in the kidneys and renal tubules of freshwater teleosts such as the smallmouthed bass (Micropterus dolomieu) and the lake minnow (Notropis sp.) (Jampol and Epstein, 1970).

Saltwater teleosts maintain a serum sodium level of about 180 - 200 mEq / liter (Jampol and Epstein, 1970). Water lost by osmosis is replaced by drinking quantities of seawater varying from 7 to 40% of the body weight per day (Maetz and Skadhauge, 1968; Parry, 1966; Smith, 1930). The sodium contained in the ingested water is excreted by the gills against a concentration gradient (Parry, 1966; Maetz and Garcia - Romeu, 1964). In addition, the gills must excrete sodium that enters the body by diffusion across the gill surface (Maetz, 1964). Consistent with these ion transport requirements,  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity is high in the gills and low in the kidneys and renal tubules of seawater teleosts such as the sea raven (Hemitripterus americanus), the long - horned sculpin (Myoxociphalus octodecimspinosus), and the goosefish (Lophius americanus) (Jampol and Epstein, 1970).

Ion influx and efflux are altered when these stenohaline teleosts are transferred to a different environmental salinity, disrupting their homeostasis and resulting in death. Some teleosts, referred to as euryhaline, can survive and adapt to abrupt changes in salinity (Zaugg and McLain, 1971). Such euryhaline teleosts are of special interest because of their physiological responses to varying salinities through changes in enzymatic activity. Jampol and Epstein (1970) reported that the activity of  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase in gill filaments and intestinal mucosa doubles when the American eel (Anguilla rostrata) is adapted to seawater.



Sodium - potassium - ATPase activity increases fourfold in the gills of the Japanese eel (Anguilla japonica), the rainbow trout (Salmo gairdnerii irideus), and the goby (Acanthogobius flavimanus) when these fishes are transferred from freshwater to seawater (Jampol and Epstein, 1970). In the killifish, Fundulus heteroclitus, the ATPase activity in the gills increases during adaptation to seawater (Epstein et al., 1967), while it decreases in the kidneys (Epstein et al., 1969). Towle and Gilman (1973) also found that the  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity in the gills of F. heteroclitus increases during adaptation to 30 ppt salt water and reaches a constant level within one - half hour after transfer from freshwater.

Osmoregulation in teleosts as well as other animals during adaptation to varying salinities has been shown to involve the endocrine system. The pituitary gland seems to play an important role in the osmoregulation of teleosts in freshwater. Burden (1956) showed that hypophysectomy destroyed the ability of F. heteroclitus to survive in freshwater, and speculated that prolactin or some unknown adeno - hypophysial factor is necessary for osmoregulation in freshwater. In later investigations, it was discovered that prolactin is the freshwater survival factor missing in the hypophysectomized killifish (Pickford and Phillips, 1959). Since these investigations, mammalian prolactin (usually ovine) has been used to restore freshwater tolerance

in the following hypophysectomized teleosts: the sailfin molly (Poecilia latipinna) (Ball and Oliverreau, 1964; Pickford et al., 1966), the moonfish (Xiphophorus maculatus) (Schreibman and Kallman, 1966; Pickford et al., 1966), the mosquitofish (Gambusia sp.) (Chambolle, 1966), the cichlid (Tilapia mossambica) (Dharmamba et al., 1967; Pickford et al., 1966), and the medaka (Oryzias latipes) (Utida et al., 1971). Prolactin is essential for life in freshwater in the above species. However, after hypophysectomy, prolactin is not essential for life in the plains killifish (Fundulus kansae) (Stanley and Fleming, 1966), the European eel (Anguilla anguilla) (Maetz et al., 1967), the flounder (Platichthys flesus) (MacFarlane, 1971), and the goldfish (Carassius auratus) (Lahlou and Sawyer, 1969). Nevertheless, without prolactin, some osmoregulatory impairments occur, which may be corrected by prolactin treatment.

The pituitary - interrenal (adrenal) axis seems to play the important role in the osmoregulation of teleosts in seawater. Hypophysectomy or interrenalectomy reduces the sodium exchange rate as well as the net sodium efflux in the eel gill, and adrenocorticotrophic hormone or cortisol restores them to the normal level (Mayer et al., 1967; Maetz, 1969). Cortisol has been shown to be involved in seawater adaptation in A. anguilla, and A. japonica (Maetz, 1969; Chester - Jones et al., 1969), A. rostrata (Epstein et al., 1971; Butler and Carmichael,

1972), C. auratus (Lahlou and Giordan, 1970), and F. heteroclitus (Epstein et al., 1967; Pickford et al., 1970).

The relationship between  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity and hormonal control has not been well defined and recent investigations have attempted to study this correlation. An increased concentration of plasma cortisol during seawater adaptation was found in A. rostrata (Forrest et al., 1973), A. japonica (Hirano, 1969), and A. anguilla (Ball et al., 1971). This burst of cortisol secretion occurs shortly after the eels make their first contact with seawater and plays an important role in conditioning the animal to withstand the osmotic stress (Forrest et al., 1973). In A. rostrata, hypophysectomy prevents the increase in ATPase activity observed upon transfer of control fish to seawater; the enzyme activity level can be restored by cortisol (Forrest et al., 1973; Epstein et al., 1971; Butler and Carmichael, 1972). Cortisol injections into intact freshwater A. japonica were found to induce a rise in the branchial  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity (Kamiya, 1972). This increase in enzyme activity produced by cortisol is of special interest because it is most unlikely that transport of sodium across the gills was increased while the fish were in freshwater. Thus increased ATPase activity is not due to a secondary adaption of the enzyme to a primary increase in  $\text{Na}^+$  transport as was previously speculated (Maetz, 1969). Epstein et al. (1971) observed striking

and unexpected pigmentation changes in specimens of A. rostrata that were injected with cortisol for more than 7 to 10 days. Such fish lost their yellow color and became silver, resembling the silver hue of A. anguilla migrating to seawater.

If cortisol exerts a direct effect on the gills of the eel, then one might expect that this hormone would be bound to the gills, as found in other cases of hormones binding to target tissues. Goodman and Butler (1972) found that gills of freshwater A. rostrata do bind large amounts of injected radioactive cortisol. Towle and Gilman (1973) showed control of the ATPase activity during adaptation of F. hetero-clitus to 30 ppt salt water does not depend on synthesis of new enzyme molecules and that control must occur by activation of preexisting enzyme. On the basis of studies cited above, they speculated that cortisol may be involved in the direct regulation of ATPase activity.

Aldosterone, like cortisol, is another hormone produced in the adrenal cortex and functions as a mineralocorticoid by affecting electrolyte metabolism. Most work with aldosterone was done on in vitro toad bladder in which aldosterone was shown to increase sodium transport (Crabbe, 1961), and on in vivo rat kidney where results were controversial. It is well established that the activity of the ATPase in membrane fragments of kidney tissue is reduced after adrenalectomy (Chignell and Titus, 1966; Jorgensen, 1968; Katz and Epstein, 1967;

Landon et al., 1966). However, mineralocorticoids were found to have no effect on ATPase activity in some cases (Chignell and Titus, 1966; Landon et al., 1966), while increasing activity in other cases (Jorgensen, 1968). There is little information on aldosterone as it effects sodium transport in teleosts and the results of the investigations are of conflicting nature (Mayer and Maetz, 1967; Favre, 1960; Motais, 1967; Henderson and Chester - Jones, 1967; Holmes et al., 1963).

Increasing evidence is now available to support the hypothesis that many hormones bring about intracellular effects through the mediation of a second messenger that has been shown to be 3', 5' - cyclic adenosine monophosphate (cyclic AMP) (Sutherland et al., 1968; Turner and Bagnara, 1971; Jost and Rickenberg, 1971). Through the action of adenyl cyclase, ATP is converted to cyclic AMP, which then acts within the effector cell to produce the appropriate hormonal response. Thus, specific hormonal events can possibly be mimicked in the effector system by the application of cyclic AMP. It has been established that cyclic AMP mimics the actions of antidiuretic hormone on the permeability of the toad urinary bladder to water and sodium (Orloff and Handler, 1965). Bastide and Jard (1968) also showed that cyclic AMP mimics the action of adrenalin by increasing sodium transport across the amphibian skin.

This investigation was undertaken to determine if the in vitro effects of prolactin and cortisol on the branchial  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase activity in freshwater adapted F. heteroclitus concur with the in vivo responses found in the articles discussed above. The effects of aldosterone on ATPase activity were of interest because little work has been done using aldosterone on teleosts. Experiments with cyclic AMP were included to determine if it might be involved in the regulation of branchial ATPase activity.

## METHODS AND MATERIALS

### General

The Fundulus heteroclitus used in these experiments were adult males, collected from Mobjack Bay at Severn, Virginia. The fish were kept in an 80 - gallon Living Stream aquarium filled with tap water, which was constantly filtered and maintained at a temperature of 18 C. Overhead fluorescent laboratory lights served as the light source and were set for a 12 hour daylight period. Fish were fed Tetramin Fish Food daily. The fish were adapted to freshwater for at least two weeks before experimentation.

### Gill Microsome Preparation

Microsomes were prepared according to a modified procedure of Hendler, et al. (1972). Groups of four to six specimens were sacrificed on various days and at approximately the same time of day. After decapitation, gills were excised and rinsed in cold homogenizing solution (0.25 M sucrose, 6 mM ethylenediamine tetraacetic acid, and 20 mM imidazole pH 6.8) and blotted. The gill tissue was then removed surgically from the cartilaginous arches and pooled to give the 500 - 800 mg gill tissue weight required for preparation of the microsomal fraction. The gill preparation was maintained at 0 - 4 C throughout

the rest of the procedure. To each 100 mg of tissue, 2 ml of homogenizing solution containing 0.1% (w/v) sodium deoxycholate was added. The tissue was homogenized in a Potter - Elvehjem apparatus with a smooth Teflon pestle for 18 strokes at 1,725 rpm. The homogenate was filtered through a double layer of cheesecloth and centrifuged at  $10,800 \times g$  at 0 C for 35 minutes. The supernatant was carefully removed and centrifuged at  $105,000 \times g$  at 0 C for 60 minutes. The microsomal pellet was resuspended in 0.3 ml of homogenizing solution per 100 mg original gill weight and homogenized for three strokes at 1,725 rpm. This microsomal preparation was frozen and maintained at -20 C until used for the measurement of  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity.

#### ATPase Assay

The assay procedure of Epstein et al. (1967) as modified by Gilman (1973) was used to measure the  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase activities in the microsomal fractions of the gills of F. heteroclitus. Total ATPase activity was measured in a medium containing 20 mM imidazole - HCl (pH 7.8), 30 mM KCl, 100 mM NaCl, and 0.1 ml of the gill microsome preparation. To this mixture was added prolactin (7.5 IU) or varying concentrations ( $10^{-7}$  -  $10^{-3}$  M) of cortisol, aldosterone, or cyclic AMP. The ouabain - insensitive activity was measured in a mixture containing the same ingredients as the total ATPase activity mixture except that 130 mM NaCl and 1.0 mM ouabain were substituted for 30 mM KCl and



100 mM NaCl. All tests were run in duplicate. The tubes were incubated in a water bath at 25 C for five minutes. The reaction was then initiated by the addition of 0.2 ml of 50 mM ATP and 50 mM  $\text{MgCl}_2$  (adjusted with Tris to pH 7.0), making a final reaction volume of 2.0 ml. All tubes were vortexed. The mixture was then incubated at 25 C for exactly 30 minutes after which the reaction was stopped by the addition of 2.0 ml of 10% trichloroacetic acid. The tubes were placed in an ice bath for 10 minutes and then centrifuged at  $10,000 \times g$  at 0 C for 10 minutes. Inorganic phosphate (Pi) in the supernatant was then measured by the Fiske - SubbaRow (1925) method. Protein concentrations of the microsome preparations were measured by the Lowry et al. (1951) method. The ouabain - sensitive  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase specific activity was defined as the difference between the total enzymatic activity (without ouabain) and the  $\text{Mg}^{++}$  - dependent, ouabain - insensitive ATPase activity (with ouabain). The specific activity of the enzyme was expressed as u moles of Pi released from ATP per mg of protein per minute.

## RESULTS AND DISCUSSION

Prolactin injections have been shown to reduce sodium efflux by decreasing the activity of  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase in hypophysectomized F. heteroclitus (Pickford et al., 1970; Utida et al., 1971), A. japonica (Kamiya, 1972), and F. kansae (Fleming and Ball, 1972) which had just been transferred from seawater to freshwater. This suggests that prolactin is involved in freshwater adaptation, an idea that is widely accepted. However, the effects of prolactin on fish already adapted to freshwater are not clear. Although Stanley and Fleming (1967) found that prolactin injections had no effect on ATPase activity in intact freshwater adapted F. kansae, Fleming and Ball (1972) found that a higher dose of prolactin reduced enzyme activity in such fish which had been hypophysectomized.

In the present study, prolactin reduced branchial  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase of freshwater adapted killifish to 79% and 85% of the controls in two different experiments (Table 1). There was no change in activity in a third experiment, possibly due to the low activity already present. The results presented here compare favorably with the effects observed in vivo by Fleming and Ball (1972).

In seawater adapted F. heteroclitus, prolactin injections decrease

branchial ATPase activity (Mayer, 1970) but have no effect on enzyme activity in A. japonica (Kamiya, 1972) or A. anguilla (Mayer, 1970). The ATPase of F. heteroclitus thus appears to be more sensitive to prolactin control than the enzyme from other species.

Injections of cortisol into intact freshwater adapted A. japonica induce an increase in branchial  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity (Kamiya, 1972). Several investigators have reported similar increases in ATPase activity in hypophysectomized seawater adapted teleosts injected with cortisol (Epstein et al., 1967; Epstein et al., 1971; Pickford et al., 1970). It was thus hypothesized that cortisol might increase enzyme activity in the present in vitro studies. As presented in Table 2, cortisol produced an increase and decrease in branchial ATPase activity of freshwater adapted killifish and on one occasion had no effect. The reason for these conflicting results are unknown. In some instances, it has been reported that such varying results may occur due to differences in animal husbandry, experimental techniques, stress in handling, seasonal variations, or differences in the natural adaptation of the fish (Milne et al., 1971; Lam, 1972). In any case, the present study offers no evidence to support a direct effect of cortisol on  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity.

Administration of aldosterone has been shown to increase (Henderson and Chester - Jones, 1967) or decrease (Holmes et al., 1963)

active sodium absorption by gills of freshwater adapted eels. It was found in the present study that aldosterone decreases gill  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity of freshwater adapted killifish (Table 3), suggesting that this hormone may reduce sodium influx in vivo. Although no dose - response relationship was found, aldosterone decreased enzyme activity in every case, ranging from 95% to 50% of the control.

The concentration of aldosterone is very low (about  $10^{-8}$  M) in the circulation of some freshwater teleosts (Chavin and Singley, 1972) and has not been found in many other fishes (Bentley, 1971). In the experiments presented here, the aldosterone concentrations used ( $10^{-7}$  to  $10^{-3}$  M) may thus have been too high to give an accurate picture of the in vivo situation. Within this concentration range, however, aldosterone appears to have a direct effect on  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase activity. The only previous indication of such a direct effect was obtained by Hendler et al. (1972) with preparations from mammalian kidney.

Cyclic AMP has been shown to increase sodium transport in toad bladder (Kirchberger et al., 1971; Orloff and Handler, 1965; Sugita et al., 1973) and amphibian skin (Bastide and Jard, 1968). In addition, Riddick et al (1971) found that dibutyryl cyclic AMP increases potassium transport in duck erythrocytes. No studies have been done,

however, on the direct effect of cyclic AMP on  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase. The results reported here clearly show that cyclic AMP stimulates branchial  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase from freshwater adapted F. heteroclitus (Table 4 and Figure 1). Maximal stimulation (158% of the control) was obtained with  $10^{-3}$  M cyclic AMP, and a clear dose - response relationship was observed in two out of three experiments (Figure 1).

Cyclic AMP is known to activate several enzymatic processes involving protein phosphorylation (Jost and Rickenberg, 1971). One of the steps in the reaction mechanism of  $\text{Na}^+$ ,  $\text{K}^+$  - ATPase requires the phosphorylation of a membrane protein (Bader et al., 1968). The stimulation of ATPase activity by cyclic AMP, as reported here, is consistent with these observations.

The actions of many hormones appear to be mediated by cyclic AMP (Jost and Rickenberg, 1971). Whether cyclic AMP mediates the in vivo action of a hormone on sodium transport in the gills of F. heteroclitus requires further investigation.

#### LITERATURE CITED

- Bader, H., Post, R. L., and Bond, G. H. (1968). Comparison of sources of a phosphorylated intermediate in transport ATPase. Biochim. Biophys. Acta. 150, 41 - 46.
- Ball, J. N., Chester - Jones, I., Forster, M. E., Hargreaves, G., Hawkins, E. F., and Milne, K. P. (1971). Measurement of plasma cortisol levels in the eel Anguilla anguilla in relation to osmotic adjustments. J. Endocrinol. 50, 75 - 96.
- Ball, J. N., and Oliverreau, M. (1964). Rôle de la prolactine dans la survie en eau douce de Poecilia latipinna hypophysectomisé et arguments en faveur de sa synthèse par les cellules érythrosinophiles de l'hypophyses des téléostéens. Compt. Rend. Acad. Sci. 259, 1443 - 1446.
- Bastide, F., and Jard, S. (1968). Actions de la noradrénaline et de l'ocytocine sur le transport actif de sodium et la perméabilité a l'eau de la peau de grenouille. Rôle du 3', 5' - AMP cyclique. Biochim. Biophys. Acta (Amst). 150, 113 - 123.
- Bentley, P. J. (1971). Endocrines and Osmoregulation. Springer - Verlag, New York, pp. 1 - 88, 199 - 254.
- Burden, C. E. (1956). The failure of hypophysectomized Fundulus heteroclitus to survive in freshwater. Biol. Bull. 110, 8 - 28.

- Butler, D. G., and Carmichael, F. J. (1972). ( $\text{Na}^+ - \text{K}^+$ ) - ATPase activity in eel (Anguilla rostrata) gills in relation to changes in environmental salinity: Role of adrenocortical steroids. Gen. Comp. Endocrinol. 19, 421 - 427.
- Chambolle, P. (1966). Recherches sur l'allongement de la durée de survie après hypophysectomie chez Gambusia sp. Compt. Rend. Acad. Sci. 262, 1750 - 1753.
- Chavin, W., and Singley, J. A. (1972). Adrenocorticoids of the goldfish (Carassius auratus L.). Comp. Biochem. Physiol. 42, 547 - 562.
- Chester - Jones, I., Chan, D. K. O., Henderson, I. W., and Ball, J. N. (1969). The adrenocorticosteroids, adrenocorticotropin and the corpuscles of Stannius. In Hoar, W. S., and Randall, D. J. (1969). Fish Physiology. Vol. 2, Academic Press, New York. pp. 321 - 376.
- Chignell, C. F., and Titus, E. D. (1966). Effect of adrenal steroids on a  $\text{Na}^+$  - and  $\text{K}^+$  - requiring adenosine triphosphatase from rat kidney. J. Biol. Chem. 241, 5083 - 5089.
- Crabbe, J. (1961). Stimulation of active sodium transport across the isolated toad bladder after injection of aldosterone to the animal. Endocrinol. 69, 673 - 682.
- Dharmamba, M., Handin, R. I., Nandi, J., and Bern, H. A. (1967). Effect of prolactin on freshwater survival and on plasma osmotic pressure of hypophysectomized Tilapia mossambica. Gen. Comp. Endocrinol. 9, 295 - 302.

- Ensor, D. M., and Ball, J. N. (1972). Prolactin and osmoregulation in fishes. Fed. Proc. 31, 1615 - 1623.
- Epstein, F. H., Cynamon, M., and McKay, W. (1971). Endocrine control of Na - K - ATPase and seawater adaptation in Anguilla rostrata. Gen. Comp. Endocrinol. 16, 323 - 328.
- Epstein, F. H., Katz, A. I., and Pickford, G. E. (1967). Sodium and potassium activated adenosine triphosphatase of gills: role in adaptation of teleosts to salt water. Science. 167, 1245 - 1247.
- Epstein, F. H., Mantius, A., Weinstein, E., Katz, A. I., and Pickford, G. E. (1969). Sodium and potassium activated adenosine triphosphatase in kidneys of Fundulus heteroclitus adapted to fresh and salt water. Yale J. Biol. Med. 41, 388 - 393.
- Favre, L. C. (1960). Recherches sur l'adaptation de Cyprin dorés à des milieux de solinté diverse et sur son déterminisme endocrinien. Diplôme d'Etudes Supérieures, Paris.
- Fiske, C. H., and SubbaRow, Y. (1925). The colorimeter determination of phosphorus. J. Biol. Chem. 66, 375.
- Fleming, W. R., and Ball, J. N. (1972). The effect of prolactin and ACTH on the sodium metabolism of Fundulus kansae held in deionized water, sodium - enriched freshwater, and concentrated seawater. Z. vergl. Physiologie. 76, 125 - 134.
- Forrest, J. N. Jr., Cohen, A. D., Schon, D. A., and Epstein, F. H. (1973). Na transport and Na - K - ATPase in the gills during adaptation to seawater: effects of cortisol. Amer. J. Physiol. 224, 709 - 713.



- Forrest, J. N. Jr., MacKay, W. C., Gallagher, B., and Epstein, F. H. (1973). Plasma cortisol response to salt water adaptation in the American eel Anguilla rostrata. Amer. J. Physiol. 224, 714 - 717.
- Fraser, R. (1970). The effect of steroids on the transport of electrolytes through membranes. In Smellie, R. M. S., ed. (1971). The Biochemistry of Steroid Hormone Action. Academic Press, New York. pp. 101 - 127.
- Gilman, M. E. (1973). Changes in the ( $\text{Na}^+ + \text{K}^+$ ) - Dependent ATPase activity in gills of the euryhaline teleost, Fundulus hetero-clitus, upon adaptation to freshwater and salt water as determined by specific activities and  $^3\text{H}$  - ouabain binding sites. Masters Thesis, University of Richmond.
- Goodman, J. H., and Butler, D. G. (1972). Localization of 4 -  $^{14}\text{C}$  cortisol in the gills of the North American eel (Anguilla rostrata Leseur). Comp. Biochem. Physiol. 42A, 277 - 296.
- Harris, E. J. (1972). Transport and Accumulation in Biological Systems. Butterworths, London, and University Park Press, Baltimore. pp. 1 - 454.
- Henderson, I. W., and Chester - Jones, I. (1967). Endocrine influence on the net extrarenal fluxes of sodium and potassium in the European eel (Anguilla anguilla L.). J. Endocrinol. 37, 319 - 325.
- Hendler, E. D., Torretti, J., Kupor, L., and Epstein, F. H. (1972). Effects of adrenalectomy and hormone replacement on  $\text{Na}^+ - \text{K}^+$  - ATPase in renal tissue. Amer. J. Physiol. 222, 754 - 759.

- Hirano, T. (1969). Effects of hypophysectomy and salinity change on plasma cortisol concentration in the Japanese eel, Anguilla japonica. Endocrinol. Japon. 16, 557 - 560.
- Hirano, T., Johnson, D. W., and Bern, H. A. (1971). Water movement in isolated flounder urinary bladder and its hormonal control by prolactin. Nature. 230, 469 - 470.
- Holmes, W. N., Phillips, J. G., and Chester - Jones, I. (1963). Adrenocortical factors associated with adaptation of vertebrates to marine environments. Rec. Prog. Hormone Res. 19, 619 - 672.
- Jampol, L. M., and Epstein, F. H. (1970). Sodium - potassium - activated adenosine triphosphatase and osmotic regulation by fishes. Amer. Physiol. 218, 607 - 611.
- Johnson, D. W. (1973). Endocrine control of hydromineral balance in teleosts. Amer. Zool. 13, 799 - 818.
- Jorgensen, P. L. (1968). Regulation of the ( $\text{Na}^+ - \text{K}^+$ ) - activated ATP hydrolyzing enzyme system in rat kidney. I. The effect of adrenalectomy and the supply of sodium on the enzyme system. Biochim. Biophys. Acta. 151, 212 - 224.
- Jost, J. - P., and Rickenberg, H. V. (1971). Cyclic AMP. Ann. Rev. Biochem. 40, 741 - 774.
- Kamiya, M. (1972). Hormonal effect on Na - K - ATPase activity in the gill of Japanese eel, Anguilla japonica, with special reference to seawater adaptation. Endocrinol. Japon. 19, 489 - 493.

- Kamiya, M., and Utida, S. (1969). Sodium - potassium - activated adenosine triphosphatase activity in gills of freshwater, marine, and euryhaline teleosts. Comp. Biochem. Physiol. 31, 671 - 674.
- Katz, A. I., and Epstein, F. H. (1967). The role of sodium - potassium - activated adenosine triphosphatase in the reabsorption of sodium by the kidney. J. Clin. Invest. 46, 1999 - 2011.
- Kirchberger, M. A., Witkum, P., and Sharp, G. W. G. (1971). On the similarity of effects of aldosterone and adenosine 3', 5' - phosphate on  $\text{Na}^+$  transport and glucose metabolism in toad bladder. Biochim. Biophys. Acta. 241, 876 - 883.
- Lahlou, B., and Giordan, A. (1970). Le contrôle hormonal des échanges et de la balance de l'eau chez le Téléostéen d'eau douce Carassius auratus, intact et hypophysectomisé. Gen. Comp. Endocrinol. 14, 491 - 509.
- Lahlou, B., and Sawyer, W. H. (1969). Electrolyte balance in hypophysectomized goldfish, Carassius auratus L. Gen. Comp. Endocrinol. 12, 370 - 377.
- Lam, T. J. (1972). Prolactin and hydromineral regulation in fishes. Gen. Comp. Endocrinol. Suppl. 3, 317 - 327.
- Landon, E. J., Jasab, N., and Forte, L. (1966). Aldosterone and sodium - potassium - dependent ATPase activity of rat kidney membranes. Amer. J. Physiol. 211, 1050 - 1056.

Lowry, O. H., Fosebrough, N. S., Farr, A. L., and Randail, R. J. (1951).

Protein measurements with the folin phenol reagent. J. Biol. Chem.  
193, 265.

MacFarlane, N. Ph. D. Thesis, University of Stirling, 1971, as reported  
by Ensor and Ball, 1972.

Maetz, J. (1964). Recherches sur la perméabilité aux électrolytes de  
la branchie des poissons et sa régulation endocrinienne. Extrait  
du Bull. Inform. Sci. Tech. Commissariat à l'Energie Atomique.  
No. 86, 1 - 60.

Maetz, J. (1969). Observations on the role of the pituitary - interrenal  
axis in the ion regulation of the eel and other teleosts. Gen.  
Comp. Endocrinol. Suppl. 2, 299 - 316.

Maetz, J., and Garcia - Romeu, F. (1964). The mechanism of sodium and  
chloride uptake by the gills of a freshwater fish, Carassius  
auratus. J. Gen. Physiol. 47, 1109 - 1116.

Maetz, J., Mayer, N., and Chartier - Baraduc, M. M. (1967). La balance  
minérale du sodium chez Anguilla anguilla en eau de mer, en eau  
douce et au cours de transfert d'un milieu à l'autre; effets de  
l'hypophysectomie et de la prolactine. Gen. Comp. Endocrinol.  
8, 177 - 188.

Maetz, J., and Skadhauge, F. (1968). Drinking rates and gill ionic  
turnover in relation to external salinities in the eel. Nature.  
217, 371 - 373.

- Mayer, N. (1970). Bull. Inform. Sci. Tech. Commissariat a l'Energie Atomique. 146, 45. As reported by Ensor and Ball, 1972.
- Mayer, N., and Maetz, J. (1967). Axe hypophyse - interénalien et osmo - régulation chez l'anguille en eau de mer. Etude de l'animal intact avec notes sur les perturbations de l'équilibre minérale produites par l'effet de "choe." Compt. Rend. Acad. Sci. 264, 1632 - 1635.
- Mayer, N., Maetz, J., Chan, D. K. O., Forster, M., and Chester - Jones, I. (1967). Cortisol, an excreting factor in the eel (Anguilla anguilla L.) adapted to seawater. Nature (London). 214, 1118 - 1120.
- Milne, K. P., Ball, J. N., and Chester - Jones, I. (1971). Effects of salinity, hypophysectomy, and corticotropin on branchial  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase in the eel, Anguilla anguilla L. J. Endocrinol. 49. 177 - 178.
- Motais, R. (1967). Les mécanismes d'échanges ioniques branchiaux chez les téléostéens. Leur rôle dans l'osmorégulation. Ann. Inst. Oceanog. 5, 1 - 83.
- Orloff, J., and Handler, J. S. (1965). Cellular mode of action of antidiuretic hormone. J. Clin. Pathol. 18, 533.
- Parry, G. (1966). Osmotic adaption in fishes. Biol. Rev. 41, 392 - 444.
- Pickford, G. E., Griffith, R. W., Torretti, J., Hendler, J., and Epstein, F. H. (1970). Branchial reduction and renal stimulation of ( $\text{Na}^+$ ,  $\text{K}^+$ ) - ATPase by prolactin in hypophysectomized killifish in freshwater. Nature. 228, 378 - 379.

- Pickford, G. E., Pang, P., and Sawyer, W. H. (1966). Prolactin and serum osmolarity of hypophysectomized killifish, Fundulus heteroclitus, in freshwater. Nature. 209, 1040 - 1041.
- Pickford, G. E., and Phillips, J. G. (1959). Prolactin, a factor in survival of hypophysectomized killifish in freshwater. Science. 130, 454 - 455.
- Potts, W. T. W. (1968). Osmotic and ionic regulation. Ann. Rev. Physiol. 30, 92 - 96.
- Riddick, D. H., Kregenow, F. M., and Orloff, J. (1971). The effect of norepinephrine and dibutyryl cyclic adenosine monophosphate on cation transport in duck erythrocytes. J. Gen. Physiol. 57, 752 - 766.
- Schreibman, M. P., and Kallman, K. O. (1966). Endocrine control of freshwater tolerance in teleosts. Gen. Comp. Endocrinol. 6, 144 - 155.
- Skou, J. C. (1957). The influence of some cations on an adenosine triphosphatase from peripheral nerves. Biochim. Biophys. Acta. 12, 12 - 20.
- Skou, J. C. (1965). Enzymatic basis for active transport of  $\text{Na}^+$  and  $\text{K}^+$  across cell membranes. Physiol. Rev. 45, 596 - 619.
- Smith, H. W. (1930). The absorption and excretion of water and salts by marine teleosts. Amer. J. Physiol. 93, 480 - 505.

- Stanley, J. G., and Fleming, W. R. (1966). Effect of hypophysectomy on the function of the kidney of the euryhaline teleost, Fundulus heteroclitus. Biol. Bull. 130, 430 - 441.
- Stanley, J. G., and Fleming, W. R. (1967). Effect of prolactin and ACTH on the serum and urine levels of Fundulus kansae. Comp. Biochem. Physiol. 20, 199 - 208.
- Sugita, M., Urakabe, S., Handler, J. S., and Orloff, J. (1973). The effect of some nucleotides and related compounds on sodium transport and water permeability of the urinary bladder of the toad. Comp. Biochem. Physiol. 45A, 157 - 167.
- Sutherland, E. W., Robison, G. A., and Butcher, R. W. (1968). Some aspects of the biological role of adenosine 3', 5' monophosphate (cyclic AMP). Circulation. 37, 279 - 306.
- Towle, D. W., and Gilman, M. E. (1973).  $\text{Na}^+$ ,  $\text{K}^+$  - dependent ATPase in gills of salt - challenged killifish: activation, synthesis, or degradation? Amer. Zool. 13, 1310.
- Turner, C. D., and Bagnara, J. T. (1971). General Endocrinology. W. B. Saunders Co., Philadelphia.
- Utida, S., Hatai, S., Hirano, T., and Kamemoto, F. K. (1971). Effect of prolactin on survival and plasma sodium levels in hypophysectomized Medaka Oryzias latipes. Gen. Comp. Endocrinol. 16, 566 - 573.
- Utida, S., Hirano, T., Cide, E., Ando, M., Johnson, D. W., and Bern, H. A. (1972). Hormonal control of the intestine and urinary bladder in teleost osmoregulation. Gen. Comp. Endocrinol. Suppl. 3, 317 - 327.

Zaugg, W. S., and McLean, L. R. (1971). Gill sampling as a method of following biochemical changes: ATPase activities altered by ouabain injections and salt water adaptation. Comp. Biochem. Physiol. 38B, 501 - 506.



Table 1. Effect of prolactin (7.5 IU) on  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase activity in gill microsomes of freshwater adapted Fundulus heteroclitus.

<u>Control Activity</u> <u>(umoles Pi/mg/min)</u>	<u>Activity with Hormone</u> <u>(umoles Pi/mg/min)</u>	<u>% of Control</u>
.525	.415	79
.274	.274	100
.418	.355	85

---

Table 2. Effect of cortisol on  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase activity

in gill microsomes of freshwater adapted Fundulus heteroclitus.

<u>Experiment 1</u>	<u>Cortisol Conc (M)</u>	<u><math>\text{Na}^+</math>, <math>\text{K}^+</math> - ATPase Activity (umoles Pi/mg/min)</u>	<u>% of Control</u>
	0	.301	100
	$10^{-7}$	.167	55
	$10^{-6}$	.301	100
	$10^{-5}$	.267	89
	$10^{-4}$	.267	89
	$10^{-3}$	.267	89
<u>Experiment 2</u>			
	0	.753	100
	$10^{-7}$	1.050	139
	$10^{-6}$	.904	120
	$10^{-5}$	.602	80
	$10^{-4}$	.979	130
	$10^{-3}$	1.050	139

Table 3. Effect of aldosterone on  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase activity  
in gill microsomes of freshwater adapted Fundulus heteroclitus.


<u>Experiment 1</u>	<u>Aldosterone Conc (M)</u>	<u><math>\text{Na}^+</math>, <math>\text{K}^+</math> - ATPase Activity (<math>\mu\text{moles Pi/mg/min}</math>)</u>	<u>% of Control</u>
	0	.335	100
	$10^{-7}$	.169	50
	$10^{-6}$	.246	73
	$10^{-5}$	.231	69
	$10^{-4}$	.277	83
	$10^{-3}$	.262	78
<u>Experiment 2</u>			
	0	.726	100
	$10^{-7}$	.630	87
	$10^{-6}$	.458	63
	$10^{-5}$	.687	95
	$10^{-4}$	.687	95
	$10^{-3}$	.687	95

---

Table 4. Effect of cyclic AMP on  $\text{Na}^+$ ,  $\text{K}^+$  - activated ATPase activity  
in gill microsomes of freshwater adapted Fundulus heteroclitus.

<u>Experiment 1</u>	<u>Cyclic AMP Conc (M)</u>	<u><math>\text{Na}^+</math>, <math>\text{K}^+</math> - ATPase Activity (umoles Pi/mg/min)</u>	<u>% of Control</u>
	0	.320	100
	$10^{-7}$	.352	110
	$10^{-6}$	.192	60
	$10^{-5}$	.368	115
	$10^{-4}$	.400	125
	$10^{-3}$	.432	135
<u>Experiment 2</u>			
	0	.795	100
	$10^{-7}$	.795	100
	$10^{-6}$	.663	83
	$10^{-5}$	.596	72
	$10^{-4}$	.861	108
	$10^{-3}$	.954	120
<u>Experiment 3</u>			
	0	.428	100
	$10^{-7}$	.428	100
	$10^{-6}$	.500	117
	$10^{-5}$	.535	125
	$10^{-4}$	.642	150
	$10^{-3}$	.678	158

Figure 1. The dose - response curve for Experiments 1 (O) and 3 (X)  
on the effects of cyclic AMP on in vitro  $\text{Na}^+$ ,  $\text{K}^+$  -  
activated ATPase activity in gill microsomes of freshwater  
adapted Fundulus heteroclitus.



## VITA

Stephen Burke Kerby was born on November 11, 1949 in Washington, D. C. He attended parochial school in Arlington County, Virginia, and public schools in Loudoun County, Virginia, and was graduated from Loudoun County High School in June 1967. He received an Associate of Science degree from Northern Virginia Community College in June 1969, and his Bachelor of Science degree from George Mason University in December, 1971, with a major in biology. He entered the University of Richmond Graduate School in February, 1972. While at Richmond, he became a member of Beta Beta Beta Honorary Biological Society. He completed the requirements for the Master of Science degree from the University of Richmond in May 1974.